

Deriving Optimal Mean Indoor Concentration Threshold Levels of PM_{2.5} and VOC for Detecting Respiratory Symptoms among Pregnant Women in Ndola and Masaiti, Zambia

David Mulenga ^{a,1}, Seter Siziya ^a

^a Copperbelt University Michael Chilufya Sata School of Medicine Public Health Unit, Ndola, Zambia,

Received 19 December 2019; revised 30 January 2020; accepted 30 January 2020

Abstract

The burden of air pollution-related morbidity and mortality in developing countries will continue if limited air monitoring and lack of air quality standards continue. This paper focuses on deriving optimal mean indoor threshold concentration levels of particulate matter (PM_{2.5}) and volatile organic compounds (VOCs) for detecting respiratory symptoms among pregnant women in Ndola and Masaiti, Zambia.

The study involved 1,170 consenting pregnant women in a cross sectional study using a standard questionnaire. Lung function tests were conducted and indoor PM_{2.5} and VOCs monitored in houses.

Biomass was the main cooking fuel. Indoor air quality monitoring results during cooking and daily average were 501(411, 686) µg/m³ and 393 (303,578) µg/m³ respectively for PM_{2.5} and 340(318,360) ppb and 343(320, 363) ppb respectively for VOCs. Significant difference in the distribution of PM_{2.5} (p-value = 0.001) and VOC (p-value = 0.017) between rural and urban area were observed. Mean indoor PM_{2.5} and VOC varied significantly by cooking activity in both rural (p-value < 0.001) and urban (p-value < 0.001) areas. Similarly, fuel type for PM_{2.5} at p-value = 0.005 but no significant difference for VOC at p-value = 0.779. However, there was a significant association between mean indoor VOC and forced vital capacity (FVC). Mean indoor PM_{2.5} was significantly higher in households that presented with respiratory symptoms than those without respiratory symptoms for both rural (p-value = 0.011) and urban areas (p-value < 0.001). Exposing a pregnant woman to mean indoor PM_{2.5} of 418µg/m³ in rural areas and 372.3µg/m³ in urban areas increased the risk of having at least one respiratory symptom. Statistical significant associations were observed between mean indoor PM_{2.5} and respiratory symptoms.

Household air pollution levels in Zambian homes are high. Therefore, systematic PM and VOCs monitoring is critical in order to develop strategies and policies relating to improvement of air quality and respiratory health.

Keywords: Air pollution, Biomass, Particulate matter, Volatile organic compounds, Spirometry

1. Introduction

Currently, developing countries are increasingly experiencing a double burden of infectious and chronic diseases (Boutayeb, 2006) partly due to environmental conditions (Sclar et al. 2005; Zulu et

¹ Corresponding author. *Email addresses:* davykdn@gmail.com (D. Mulenga); ssiziya@gmail.com (S. Siziya)

- Perez-Padilla R, Perez-Guzman C, Baez-Saldana R, Torres-Cruz A Cooking with biomass stoves and tuberculosis: a case control study. *International Journal of Tuberculosis and Lung Disease*.2001; 5:1–7
- Pope CA 3rd, Dockery DW. Health effects of fine particulate air pollution: lines that connect. *J Air Waste Manag Assoc* 2006;56:709–42.
- Romieu I, Riojas-Rodriguez H, Marron-Mares AT, Schilman A, Perez-Padilla R, Masera O. Improved biomass stove intervention in rural Mexico: impact on the respiratory health of women. *Int J Respir Crit Care Med* 2009;180:649–56.
- Rumchev K, Spickett JT, Brown HL, et al. Indoor air pollution from biomass combustion and respiratory symptoms of women and children in a Zimbabwean village. *Indoor Air* 2007;17:468–74.
- Tielsch JM, Katz J, Zeger SL, Khatri SK, Shrestha L, Breysse P, Checkley W, Mullany LC, LeClerq SC. Designs of two randomized, community-based trials to assess the impact of alternative cookstove installation on respiratory illness among young children and reproductive outcomes in rural Nepal. *BMC Public Health* 2014, 14:1271
- Sclar ED, Garau P, Carolini G. The 21st Century health challenge of slums and cities. *Lancet*. 2005; 365(9462):901-903
- St. Helen G., Aguilera-Villalobos M, Cassidy B.E., Blount B.C., Bayer C., Hall D.B., Needham L.L., Hendry R.J., Naeher L.P., 2011. Characterization of volatile organic compounds (VOCs) Exposure from cooking fuels among a cohort of pregnant women in Trujillo, Peru. *Am J Respir Crit Care Med* 183;2011:A3894
- Thomas P, Zelikoff J (1999) Air pollutants: moderators of pulmonary host resistance against infection. In: *Air pollution and health*. Holgate ST et al., eds. Academic Press, San Diego, CA.
- WHO. Air quality guidelines, global update. <http://www.euro.who.int/Document/E90038.pdf> 2006. Accessed April 2018.
- World Health Organisation, WHO IAQ guidelines 2014; Household fuel combustion. WHO Geneva.
- Zelikoff J (1994) Woodsmoke emissions: effects on host pulmonary immune defence. *CIAR Currents*, 1:3.
- Zulu EM, Beguy D, Ezech AC, Bocquier P, Madise NJ, Cleland J, Falkingham J. Overview of migration, poverty and health dynamics in Nairobi City's slum settlements. *Urban Health: Bull NY Acad Med*. 2011; 28(Suppl 2):S185-S199